

Breast Oncology

Capecitabine

This leaflet is given in addition to your consent form(s) and booklet, 'Chemotherapy - a guide for patients and their carers'.

Your treatment

Capecitabine tablets are taken orally twice a day (10 to 12 hours apart) on days 1 to 14, followed by a 7 day break. Treatment is given every 21 days (1 cycle). Treatment may be continued until disease progression or unacceptable toxicity. You will be required to have a blood test prior to starting capecitabine to check you have the enzymes required to break down this type of chemotherapy.

If you vomit after taking a dose, do not repeat the dose. Take your next dose at your normal time.

You will have routine blood tests while on treatment as advised by your medical team. The frequency of these blood tests may decrease once you are established on treatment.

Occasionally we may not be able to go ahead with your treatment until your blood counts are back to a safe level. If this happens, your treatment may be delayed a week or dose reduced.

You must never take your treatment unless you have had your blood checked and been given the go-ahead by the doctor, nurse clinician or pharmacist.

This treatment can have serious or possibly life-threatening side effects. It is very important that you report side effects straight away.

You are also vulnerable to infection while you are having chemotherapy. Minor infections can become life-threatening in a matter of hours if left untreated. Symptoms of infection include fever, shivering, sweats, sore throat, diarrhoea, discomfort when you pass urine, cough or breathlessness. We recommend that you use a digital thermometer so you can check your temperature. You can buy one from your local chemist.

If you feel unwell, you have symptoms of an infection or your temperature is 37.5°C or above, or below 36°C, contact The Christie Hotline straight away on 0161 446 3658 (open 24 hours a day, 7 days a week).



Possible side effects

See the Cancer Research UK consent form.

You should have been given a copy of this consent form before you started treatment but it can also be found by visiting www.cancerresearchuk.org and searching 'Consent forms for SACT (Systemic Anti-Cancer Therapy)'.

See 'Chemotherapy - a guide for patients and their carers'.

You should have been given a copy of this booklet. You can also see this on The Christie website, visit www.christie.nhs.uk.

Contacts

If you have any general questions or concerns about your treatment, please ring the service under which you are being treated.

Advanced breast cancer treatment (MBC):

Please contact your consultant's secretary or your breast care nurse.

If you need information in a different format, such as easy read, large print, BSL, braille, email, SMS text or other communication support, please tell your ward or clinic nurse.

The Christie is committed to producing high quality, evidence based information for patients. Our patient information adheres to the principles and quality statements of the Information Standard. If you would like to have details about the sources used please contact the-christie.patient.information@nhs.net

For information and advice visit the cancer information centres at Withington, Oldham, Salford or Macclesfield. Opening times can vary, please check before making a special journey.



Contact The Christie Hotline for
urgent support and specialist advice
The Christie Hotline: 0161 446 3658
Open 24 hours a day, 7 days a week